

True (A) or false (B):**Puerperas' knowledge of postnatal exercises**

1. It is possible that postpartum exercises may reduce postpartum complications such as haemorrhage and sepsis.

Knowledge, attitudes, and practice in emergency contraception

2. Globally there were an estimated 42 million induced abortions in 2001, of which 48% were unsafe.
3. The World Health Organization estimated that between 2003 and 2008 the number of unsafe abortions in the world increased, with the highest rates being in eastern and central Africa.
4. A review of 1 077 articles from 113 newspapers published in 2005 demonstrated a confusion between emergency contraception and medical abortion in almost 45% of those articles.

Abnormal umbilical Doppler and placental histology in fetal growth restriction

5. Maternal underperfusion lesions include villous infarction, villous agglutination, syncytial knots and intervillous fibrin deposition.
6. In the study included in this journal, abnormal Doppler had a sensitivity of 65% and specificity of 80% for abnormal placental pathology.

Preventive role of low-molecular-weight heparin in early pregnancy loss

7. There are varying definitions of recurrent pregnancy loss, which include two or three pregnancy losses.
8. The risk of miscarriage increases to up to 30% after two consecutive early pregnancy losses.
9. Recurrent pregnancy loss remains unexplained in up to 50% of couples investigated.
10. In the study included in this journal, subcutaneous enoxaparin in a dose of 40 mg once daily improved the chance of live births in non-thrombophilic women with unexplained recurrent pregnancy loss when compared with a placebo.

Galactagogue use and breastfeeding behaviour

11. The global rate of exclusive breastfeeding in the first six months of life is reported as 38%.
12. A recent Cochrane review concluded that the use of galactagogues was associated with a significant improvement in longer term outcomes of breastfeeding in preterm infants.
13. Galactagogues are dopamine agonists such as antipsychotics, antiemetics and natural supplements which enhance lactation by increasing serum prolactin.
14. In high-income settings, galactagogue use has been reported in over 30% of parturient mothers.

Beliefs and practices in using misoprostol for induction of labour in Zimbabwe

15. A global survey published by the WHO in 2014 stated that induction of labour was performed in 9.6% of all births.
16. There is currently a clear international definition of failed induction of labour.

Embolisation of pseudoaneurysm of the uterine artery

17. Secondary postpartum haemorrhage is defined as haemorrhage occurring 6 hours to 6 weeks postpartum.

Bilateral synchronous benign ovarian neoplasms

18. The most common germ cell tumours of the ovary are benign mature cystic teratomas.
19. Benign mature cystic teratomas of the ovary are composed of endoderm and ectoderm, but not mesoderm.
20. Metastases to the ovary constitute 5% of ovarian neoplasms.

The CPD programme for SAJOG is administered by Medical Practice Consulting:

CPD questionnaires must be completed online at www.mpconsulting.co.za

A maximum of 3 CEUs will be awarded per correctly completed test.

Accreditation number: MDB015/173/02/2017 (Clinical)