

CPD QUESTIONNAIRE

1. True (A) or false (B) – click on the correct answer:
Up to 1 in 6 or more breast cancers are over-diagnosed and over-treated in routine screening mammography programmes.
2. True (A) or false (B) – click on the correct answer:
In the UK death rates from breast cancer over the last 20 years have fallen from above 40 per 100 000 women to below 30 per 100 000.
3. True (A) or false (B) – click on the correct answer:
Human papillomavirus vaccines are effective and safe in women in the age group 25 - 45 years.
4. True (A) or false (B) – click on the correct answer:
Pelvic floor exercises during pregnancy reduce the use of episiotomies and the incidence of perineal tears.
5. True (A) or false (B) – click on the correct answer:
A woman who has pre-eclampsia in her first pregnancy has a 15% chance of its recurrence in her second pregnancy.
6. True (A) or false (B) – click on the correct answer:
A woman who has pre-eclampsia in her first and second pregnancies also has a 15% chance of its recurrence in her third pregnancy.
7. True (A) or false (B) – click on the correct answer:
Cell-free fetal nucleic acid fragments are found in maternal serum from early pregnancy.
8. True (A) or false (B) – click on the correct answer:
The prevalence of Down syndrome for 30-year-old women is approximately 1 in 1 000.
9. True (A) or false (B) – click on the correct answer:
The prevalence of non-chromosomal congenital anomalies rises with age in the same way as that of chromosomal abnormalities.
10. True (A) or false (B) – click on the correct answer:
When a patient has urinary stress incontinence and prolapse, the two conditions are best dealt with as separate surgical procedures.
11. True (A) or false (B) – click on the correct answer:
Deaths within the first 24 hours of life constitute one-third of all neonatal deaths.
12. True (A) or false (B) – click on the correct answer:
The overall perinatal mortality rate for South Africa is approximately 40 per 1 000 births.
13. True (A) or false (B) – click on the correct answer:
A woman's lifetime risk of surgery for urinary incontinence or pelvic organ prolapse is approximately 10%.
14. True (A) or false (B) – click on the correct answer:
The vast majority of operations for urinary incontinence or pelvic organ prolapse are covered by prophylactic antibiotics.
15. True (A) or false (B) – click on the correct answer:
The incidence of shoulder dystocia is increasing worldwide.
16. True (A) or false (B) – click on the correct answer:
Approximately 20% of babies delivered with shoulder dystocia will suffer some sort of injury.
17. True (A) or false (B) – click on the correct answer:
Prophylactic caesarean section use is set to reduce the burden of disease caused by shoulder dystocia.
18. True (A) or false (B) – click on the correct answer:
McRoberts manoeuvre should usually be attempted as the first move in the management of shoulder dystocia.
19. True (A) or false (B) – click on the correct answer:
In the last Maternal Mortality report nearly 40% of maternal deaths in South Africa had clearly avoidable causes.
20. True (A) or false (B) – click on the correct answer:
Intrapartum asphyxia and birth trauma as a combination were the commonest cause of avoidable perinatal deaths in South Africa according to the most recent survey.

CPD questionnaires must be completed online via www.cpdjournals.org.za

After submission you can check the answers and print your certificate.

Questions may be answered up to 6 months after publication of each issue.

Accreditation number: MDB001/018/01/2009