

CPD QUESTIONNAIRE

1. True (A) or false (B) – click on the correct answer:
Venous thrombo-embolism is much more common in pregnancy than in non-pregnant women of comparable age.
2. True (A) or false (B) – click on the correct answer:
Most deep-vein thromboses in pregnancy are left-sided.
3. True (A) or false (B) – click on the correct answer:
The peak incidence of postoperative venous thrombo-embolism is intra-operative or 1 week after the operation.
4. True (A) or false (B) – click on the correct answer:
Research has shown that there is no increased risk of venous thrombo-embolism 4 weeks after abdominal surgery.
5. True (A) or false (B) – click on the correct answer:
Povidone iodine is the most widely used pre-operative skin prep in the USA.
6. True (A) or false (B) – click on the correct answer:
Povidone iodine is a better pre-operative skin prep than chlorhexidine.
7. True (A) or false (B) – click on the correct answer:
Administration of the human papillomavirus vaccines during pregnancy is associated with an increased risk of sub-optimal outcomes in terms of perinatal mortality rates.
8. True (A) or false (B) – click on the correct answer:
The human papillomavirus vaccines appear safe in pregnancy in that the rates of congenital abnormalities are no higher in exposed women than in non-exposed women.
9. True (A) or false (B) – click on the correct answer:
Women using postmenopausal hormone therapy have a lowered risk of endometrial cancer if they use progestins continuously than if they use progestins sequentially.
10. True (A) or false (B) – click on the correct answer:
The selective serotonin re-uptake inhibitor antidepressants paroxetine and fluoxetine interfere with the metabolism of tamoxifen, markedly reducing its benefits.
11. True (A) or false (B) – click on the correct answer:
Being exposed to tobacco smoke *in utero* can lead to childhood behavioural problems.
12. True (A) or false (B) – click on the correct answer:
Lung cancer sufferers have a better prognosis if they quit smoking.
13. True (A) or false (B) – click on the correct answer:
It is not necessary for lesbians to have cervical cancer cytological screening.
14. True (A) or false (B) – click on the correct answer:
Cytology screening should start at sexual debut.
15. True (A) or false (B) – click on the correct answer:
Human papillomavirus DNA testing is a useful screening test in teenage women who are sexually active.
16. True (A) or false (B) – click on the correct answer:
A bolus dose of oxytocin at caesarean section has a marked effect on mean arterial blood pressure.
17. True (A) or false (B) – click on the correct answer:
Oxytocin is antidiuretic if given in large doses.
18. True (A) or false (B) – click on the correct answer:
The menopause transition often coincides with disadvantageous changes in a woman's lipid profile.
19. True (A) or false (B) – click on the correct answer:
In postmenopausal women undergoing hysteroscopic procedures, pre-operative misoprostol after oestrogenic preparation facilitates cervical dilatation.
20. True (A) or false (B) – click on the correct answer:
Vaginal cosmetic surgery has been shown to be beneficial to women with sexual dysfunction.

CPD questionnaires must be completed online via www.cpdjournals.org.za
After submission you can check the answers and print your certificate.
Questions may be answered up to 6 months after publication of each issue.

Accreditation number: MDB001/004/01/2010