

of the world. In developing countries such as South Africa the incidences of cervical and vulvar cancer in young females has increased in the past decade. The dilemma is that most of our patients present in locally advanced stage of disease with massive tumour bulk, poor nutritional status and very low CD4 counts (usually < 200/ $\mu$ l). According to one South African survey<sup>12</sup> about 70% of patients first seek health care from traditional healers and only come to hospitals when the disease is at an advanced stage and the symptoms are unbearable. At this stage surgery is not possible and in certain cases radical chemotherapy and/or radiation cannot be offered owing to very low CD4 counts and poor general condition and performance status. In this situation palliative radiation can be delivered, encompassing the local disease with the palliative dose of 20 Gy in 5 fractions (4 Gy daily dose) or 30 Gy in 10 fractions (3 Gy daily). We have observed very good palliative results in locally advanced vulvar cancer in HIV-positive patients. This good response is probably due to clinicopathological differences in these cases and the relation to HPV with warty/basaloid features and lesser invasiveness.

The case reported here reinforces the importance of a very high suspicion of malignancy when warty vulvar lesions are seen in immunosuppressed patients. Such


lesions should be biopsied and histological evaluation should be done to detect the cancer early so that radical treatment can be offered. Furthermore, there is a need to determine the most effective doses of radiation for this cohort of patients.

A shorter version of this article has appeared in the *South African Medical Journal* (2006, **96**: 1044).

1. Serraino D, Carrieri P, Pradier C. Risk of invasive cervical cancer among women with, or at risk for, HIV infection. *Int J Cancer* 1999; **82**: 334-337.
2. Serraino D. The spectrum of AIDS-associated cancers in Africa. *AIDS* 1999; **13**: 2589-2590.
3. Carter J, Carlson J, Fowler J. Invasive vulvar tumors in young women – a disease of the immunosuppressed? *Gynecol Oncol* 1993; **51**: 307-310.
4. Kuhn L, Sun X-W, Wright TC. Human immunodeficiency virus infection and female lower genital tract malignancy. *Curr Opin Obstet Gynecol* 1999; **11**: 35-39.
5. Crum CP. Carcinoma of the vulva: epidemiology and pathogenesis. *Obstet Gynecol* 1992; **79**: 448-454.
6. Van Hoeven KH, Kovatich AJ. Immunohistochemical staining for proliferating cell nuclear antigen, Bcl2 and Ki-67 in vulval tissue. *Gynecol Pathol* 1996; **15**: 10-16.
7. Al Ghamdi A, Freedman D, Miller D, et al. Vulvar squamous cell carcinoma in young women: A clinicopathologic study of 21 cases. *Gynecol Oncol* 2002; **84**: 94-101.
8. Piura B, Rabinovich A, Shaco-Levy R, Sukenik S. Vulvar invasive squamous cell carcinoma occurring in young women with systemic lupus erythematosus. *Eur J Gynaecol Oncol* 2005; **26**(1): 103-105.
9. Park K, Monk BJ, Wilczynski S, Ito JI jun, Vasilev SA. Idiopathic CD4+ T-lymphocytopenia and recurrent vulvar intraepithelial neoplasia. *Obstet Gynecol* 1994; **84**(4 Pt 2): 712-714.
10. Regan MA, Rosenzweig BA. Vulvar carcinoma in pregnancy: a case report and literature review. *Am J Perinatol* 1993; **10**(4): 334-335.
11. Ogunleye D, Lewin SN, Huettner P, Herzog TJ. Recurrent vulvar carcinoma in pregnancy. *Gynecol Oncol* 2004; **95**: 400-401.
12. Bateman C. HIV training of traditional healers limps along. *S Afr Med J* 2004; **94**: 804-806.

## CASE REPORT

# Uterine artery embolisation in a Jehovah's Witness patient



**Peter de Jong, MB ChB, MMed, FCOG, FRCOG**  
University of Cape Town and Groote Schuur Hospital, Cape Town

**Malika van der Schyff, MB ChB, FCOG**  
Constantiaberg Hospital, Cape Town

A 30-year-old nulliparous woman was referred with life-threatening massive vaginal bleeding and failed medical treatment. She was of the Jehovah's Witness faith and refused transfusion with any blood products. Uterine artery embolisation arrested the haemorrhage.

## Presentation

The patient presented with a history of heavy vaginal bleeding with associated clots. This had necessitated the use of 20 sanitary towels per day with double padding, flooding and soiling. There was also associated dizziness, weakness and malaise. She was not sexually active, and was a devout Jehovah's Witness.

On examination she was calm, co-operative and undistressed. She was obese (body mass index 30) with no signs of hirsutism, acne or acanthosis nigricans. Her blood pressure was 120/65 mmHg with an associated tachycardia of 102/min. The peripheral oxygen saturation was 98% and fingerprick haemoglobin 2 g/dl. Her temperature was 37.4°C and no abnormalities were detected on dipstick urinalysis. She had a normal

jugular venous pressure and precordium with normal heart sounds on auscultation. She was awake, alert, well orientated and responsive to questions.

The abdomen was obese with no surgical scars present. On palpation it was soft and non-tender, and no organomegaly was detected.

The vulva and vagina were normal with normal hair distribution. The cervix appeared normal on speculum examination. The uterus was of normal size, axial and mobile. Both adnexae were non-tender and no lesions were palpated on digital examination.

Over the following 24 hours management of the patient included the following medication:

- tranexamic acid (Cyclokapron) – 1 g intravenously 8-hourly
- ibuprofen – 400 mg orally 8-hourly
- medroxyprogesterone acetate (Depo-Provera) – 300 mg intramuscular injection
- norethisterone (Primolut N) – 10 mg orally 8-hourly
- conjugated oestrogen (Premarin) – 25 mg intravenously
- antibiotics – ampicillin and metronidazole intravenously.

The above treatment did not succeed in reducing the vaginal blood loss. Further investigation revealed negative serum beta-human chorionic gonadotrophin ( $\beta$ -hCG), a normal pelvic ultrasound scan, normal thyroid function and a normal haematological profile other than continuing anaemia. Over the following 36 hours the patient's condition deteriorated. Her blood loss continued, her haemoglobin continued to drop and she became shocked. Despite extensive counselling, she continued to decline blood transfusion.

She was transferred to Groote Schuur Hospital for uterine artery embolisation (UAE) since all other options to control her bleeding had failed. She was resuscitated with crystalloids (modified Ringer's lactate) and colloids (Haesteril). Bilateral uterine artery embolisation was undertaken by an interventional radiologist. The internal iliac artery was entered through the right groin area via a #5F sheath. Both uterine arteries were embolised using gelatin (Spongistan) pledgets.

After embolisation the vaginal bleeding ceased. Resuscitation continued with crystalloids and colloids. The patient was also given intravenous erythropoietin (10 000 U intravenously once daily) and oral haematinics (ferrous sulphate 200 mg orally twice daily and folic acid 5 mg orally daily), resulting in an increase in her haemoglobin concentration from 2.5 g/dl to a discharge level of 8.9 g/dl over a period of 26 days.

The prolonged period of hypovolaemia resulted in small cerebral infarcts. A CT scan revealed small infarcts including two small hypoechoic areas adjacent to the anterior horn of the left and right lateral ventricle. She was mobilised with parallel bars and was ultimately able to walk aided by a walking cane.

She was discharged 33 days after admission on the combined oral contraceptive levonorgestrel/oestrogen (Nordette) and haematinics. She continued to need a walking cane, and initially did not return to work and was cared for by relatives. At the time of writing she has resumed her previous position as a domestic worker and is in a stable relationship.

## Discussion

UAE entails the occlusion of arterial uterine blood flow with embolisation material. It incorporates angiography to visualise the uterine arteries and is performed by an experienced interventional radiologist.

UAE been used successfully in the treatment of obstetric and gynaecological haemorrhage. It has been shown to be an effective alternative to surgical intervention in the treatment of uterine bleeding resulting from a variety of conditions, including postpartum haemorrhage, postoperative haemorrhage, cervical pregnancy, arteriovenous malformation and most recently uterine leiomyomas.<sup>1-3</sup>

UAE has had success rates of 86 - 90% in treatment of excessive menstrual blood loss, with symptomatic relief seen as early as the first menstrual cycle in the majority of patients.<sup>1</sup> In the above case, where medical management failed and surgical intervention was not a viable option, UAE proved successful in controlling blood loss. To date there have been no reports of this procedure being utilised to control haemorrhage in patients of the Jehovah's Witness faith.

After UAE, uterine viability is sustained due to a non-collateral arterial blood supply with contributions from the ovarian arteries.<sup>4</sup> Embolisation of uterine arteries has a distinct effect on leiomyomas as opposed to the myometrium, being able to stop the blood supply to leiomyomas while collateral circulation sustains the viability of the uterine tissue.<sup>1,4</sup> Embolisation involves the injection of one of three types of embolisation materials (polyvinyl alcohol micro particles/PVA, gelatin sponges or calibrated microspheres) through a catheter.<sup>5</sup> If the procedure is done electively, patients are discharged within 24 hours.<sup>1,4</sup>

## Conclusion

Uterine artery embolisation is a novel technique that can be utilised in both obstetric and gynaecological settings as a minimally invasive procedure. There are no documented cases of UAE used in a Jehovah's Witness patient in the setting of life-threatening haemorrhage.

1. Goodwin SC, Wong GCH. Uterine artery embolization for uterine fibroids: A radiologist's perspective. *Clin Obstet Gynecol* 2001; **44**(2): 412-424.
2. Florindon C, Lund N, Grove Thomsen S. Alternative treatment for symptomatic fibroids. *Curr Opin Obstet Gynecol* 2001; **3**: 491-495.
3. Broder MS, Landow WJ, Goodwin SC, Brook SC, Sherbourne CD, Harris K. An agenda for research into uterine artery embolization: Results of an expert panel conference. *J Vasc Interv Radiol* 2000; **11**: 509-515.
4. Vedantham S, Goodwin SC, McClucas B, Mohr G. Uterine artery embolization: An underused method of controlling pelvic hemorrhage. *AJOG* 1997; **176**: 938-948.
5. Worthington-Kirsch RL. Technical failure. *J Vasc Interv Radiol* 2001; **8**: Suppl 12, 256.